



## King's Research Portal

DOI:

[10.1371/journal.pone.0137514](https://doi.org/10.1371/journal.pone.0137514)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Kim, Y. R., Eom, J. S., Yang, J. W., Kang, J., & Treasure, J. (2015). The impact of oxytocin on food intake and emotion recognition in patients with eating disorders: A double blind single dose within-subject cross-over design. *PLoS One*, 10(9), [e0137514]. <https://doi.org/10.1371/journal.pone.0137514>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

# The Impact of Oxytocin on Food Intake and Emotion Recognition in Patients with Eating Disorders: A Double Blind Single Dose Within-Subject Cross-Over Design

Youl-Ri Kim<sup>1,2\*</sup>, Jin-Sup Eom<sup>3</sup>, Jae-Won Yang<sup>4</sup>, Jiwon Kang<sup>2</sup>, Janet Treasure<sup>5</sup>

**1** Department of Neuropsychiatry, Seoul Paik Hospital, Inje University, Seoul, South Korea, **2** Institute of Eating Disorders and Mental Health, Inje University, Seoul, South Korea, **3** Department of Psychology, Chungbuk National University, Cheongju, South Korea, **4** Department of Psychology, The Catholic University of Korea, Bucheon, South Korea, **5** Section of Eating Disorders, Department of Psychological Medicine, King's College London, Institute of Psychiatry, London, United Kingdom

\* [youlri.kim@paik.ac.kr](mailto:youlri.kim@paik.ac.kr)



## OPEN ACCESS

**Citation:** Kim Y-R, Eom J-S, Yang J-W, Kang J, Treasure J (2015) The Impact of Oxytocin on Food Intake and Emotion Recognition in Patients with Eating Disorders: A Double Blind Single Dose Within-Subject Cross-Over Design. PLoS ONE 10(9): e0137514. doi:10.1371/journal.pone.0137514

**Editor:** Susana Jiménez-Murcia, University Hospital of Bellvitge-IDIBELL; CIBER Fisiopatología Obesidad y Nutrición (CIBEROBN), Instituto Salud Carlos III; Department of Clinical Sciences, School of Medicine, University of Barcelona, Spain, SPAIN

**Received:** April 10, 2015

**Accepted:** August 17, 2015

**Published:** September 24, 2015

**Copyright:** © 2015 Kim et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was supported by a donation from the parents of a Korean patient with anorexia nervosa. This work was also supported by the Swiss Anorexia Nervosa Foundation. Janet Treasure receives salary support from the NIHR Biomedical Research Centre for the Institute of Psychiatry, King's College London. The funders had no role in the study

## Abstract

### Background and Aim

Social difficulties and problems related to eating behaviour are common features of both anorexia nervosa (AN) and bulimia nervosa (BN). The aim of this study was to examine the impact of intranasal oxytocin on consummatory behaviour and emotional recognition in patients with AN and BN in comparison to healthy controls.

### Materials

A total of 102 women, including 35 patients with anorexia nervosa (AN), 34 patients with bulimia nervosa (BN), and 33 healthy university students of comparable age and intelligence, participated in a double-blind, single dose placebo-controlled cross-over study. A single dose of intranasal administration of oxytocin (40 IU) (or a placebo) was followed by an emotional recognition task and an apple juice drink. Food intake was then recorded for 24 hours post-test.

### Results

Oxytocin produced no significant change in appetite in the acute or 24 hours free living settings in healthy controls, whereas there was a decrease in calorie consumption over 24 hours in patients with BN. Oxytocin produced a small increase in emotion recognition sensitivity in healthy controls and in patients with BN. In patients with AN, oxytocin had no effect on emotion recognition sensitivity or on consummatory behaviour.

### Conclusions

The impact of oxytocin on appetite and social cognition varied between people with AN and BN. A single dose of intranasal oxytocin decreased caloric intake over 24 hours in people

design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

with BN. People with BN showed enhanced emotional sensitivity under oxytocin condition similar to healthy controls. Those effects of oxytocin were not found in patients with AN.

## Trial Registration

ClinicalTrials.gov [KCT0000716](https://clinicaltrials.gov/ct2/show/study?term=KCT0000716)

## Introduction

Oxytocin is now recognised as having a central role in the neural circuits involved in social behaviour, appetite, anxiety, and stress [1,2]. These features are also characteristic of people with an eating disorder, which raises the possibility that oxytocin is involved in the pathophysiology of the disorder [3,4]. During the starvation phase of anorexia nervosa (AN), the levels of oxytocin in the cerebrospinal fluid are decreased [5–7]. There are changes in the release of oxytocin in response to a meal in both the acute illness and post-recovery conditions [8]. Moreover, an intranasal administration of oxytocin produces changes in the attentional processing of cues that are salient for people with AN. For example, the vigilance toward food and body image stimuli is reduced [9]. In addition, the attentional bias toward negative facial emotions (disgust, anger) was modified by oxytocin in AN [10]. These findings suggest that oxytocin systems may be involved in fear-related stimuli and social cue processing in AN patients.

Meanwhile, fewer studies have investigated the oxytocin system in bulimia nervosa (BN) or in binge eating disorders although some of the features of these conditions also suggest a possible dysfunction in oxytocin systems. Oxytocin has been demonstrated to be an important peptide for body weight regulation [11,12]. Animal studies suggest that oxytocin is involved in weight control [13,14] and is particularly involved in inhibiting the appetite for sugar and carbohydrates [11,15]. Oxytocin receptor antagonist injections in wild-type mice produced a preference for sucrose over fat [16]. Oxytocin receptor knockout animals consume greater amounts of sweet solutions than wild types [17] and develop late onset obesity [18–20]. Animals engineered not to express oxytocin overconsume sweetened food [21] and carbohydrates [22]. Oxytocin expression was down regulated upon long-term intermittent exposure to sugar, which may represent a form of neural adaptation to a high sugar diet [23]. Mice with dietary-induced obesity exhibit functional abnormalities in the oxytocin systems. In humans, the administration of oxytocin in obese men produced weight loss [24]. Moreover, a recent study on healthy men found that oxytocin reduced the intake of high sugar snacks [25,26]. To our knowledge, there has been no study investigating oxytocin function in patients with BN or binge eating disorders.

In addition to playing a key role in appetite, oxytocin is involved in social and emotional processing. Problems that involve social emotional cognition may underpin some of the interpersonal difficulties in patients with eating disorders, such as a reduction in emotional expression in patients with AN and problems with trust and conflict in patients with BN, which are thought to contribute to the maintenance of the disorders [27]. Social cognition is complex and includes various subprocesses. A recent meta-analysis of the various components of social cognition in patients with eating disorders found an impairment in most of the domains of social functioning [28]. Emotional recognition was impaired in patients with eating disorders but with a large variation observed between tasks, and the weakest effects were observed for static photographs of faces. To our knowledge, a more ecologically valid technique to measure the sensitivity for emotional discrimination with dynamically changing facial expressions has

yet to be used to examine eating disorders. In this study, we used this experimental paradigm to compare the emotional recognition sensitivity between people with AN or BN and healthy controls following a single dose of placebo or oxytocin.

The aim of this study was to examine the impact of an intranasal oxytocin challenge test on two possible maintaining factors of eating disorders: problems in appetite control and social cognition. The first hypothesis was that oxytocin would decrease food consumption. The second hypothesis was that social cognition would be improved in the oxytocin condition in patients with eating disorders.

## Materials and Methods

### Participants

The protocol for this study was approved by both the Institutional Review Board of the Korean Food and Drug Association (12061) and the Institutional Review Board of the Seoul Paik Hospital (IIT-2012-096) on September 16<sup>th</sup>, 2012 (See [S1–S4](#) Texts). Although this is an experimental proof of concept study, and we have established with the regulatory agencies in the UK that it does not need to be registered, we have found that other experts and journals find this confusing. Therefore, we have registered the study in the clinical trial registry [the Clinical Research Information Service (<http://cris.nih.go.kr>) (registration number: KCT0000716)]. All ongoing and related trials for this drug/intervention were registered. All of the participants provided written informed consent.

One hundred and two women (35 patients with AN, 34 patients with BN, and 33 healthy university students) completed this double-blind, placebo-controlled cross-over study. The first participant entered the study on April 4<sup>th</sup>, 2013, and the last participant completed follow-up on August 14<sup>th</sup>, 2014. The patients with AN were recruited from the inpatient ward and the patients with BN were recruited from the outpatient clinic of the Eating Disorders Clinic of the Seoul Paik Hospital in Seoul, South Korea. The diagnosis of eating disorder was confirmed by the Structured Clinical Interview from the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition [29]. All patients satisfied the DSM-5 diagnostic criteria of AN or BN. The exclusion criteria for comorbidities included active substance use disorder, psychotic disorder (schizophrenia, schizoaffective, psychosis not otherwise specified), and autism spectrum disorder. Patients taking psychiatric medications other than fluoxetine were also excluded.

The healthy controls were undergraduate or graduate students of a comparable age and intelligence who responded to an advertisement posted in the psychology department at a women's university in Seoul, South Korea. The inclusion criteria were healthy females without a history of medical or psychiatric illness and a minimum age of 17. We screened all healthy participants for eating disorders with the SCOFF questionnaire [30]. The exclusion criteria for the healthy controls included a self-reported history of major depression, bipolar, panic, or psychotic disorders, substance dependence, epilepsy, eating disorder, autism spectrum disorder, traumatic brain injury, and taking medications (including contraceptives). We also excluded the participants who were smokers, homosexual, or parous because these conditions may interfere with oxytocin functioning.

The healthy controls and patients with BN who had their period were tested during the follicular phase of their menstrual cycle (approximately days 3 through 12). None of the patients with AN were menstruating. The participants were compensated for their travelling expenses and time.


The degree of the eating psychopathology for all participants (weight, shape, eating concerns, and dietary restraint) was evaluated using the Korean version of the Eating Disorder Examination self-report version Questionnaire (EDE-Q) [31]. Depression and anxiety were

assessed for each subject by using the standardized Korean versions of the Beck Depression Inventory (BDI) [32] and the Spielberger State and Trait Anxiety Inventory (STAI) [33], respectively. The Korean version of the Ward 7-subtest short form of the Wechsler Adult Intelligence Scale [34] was used to measure the IQ of all participants.

A consort diagram is shown in Fig 1 (See S5 Text).

### Oxytocin preparation

The intranasal oxytocin spray was formulated by JW Pharmaceuticals (Seoul, South Korea) from oxytocin powder (Hemmo Pharmaceuticals, Mumbai, India). The solution was prepared by combining 35.2 mg of oxytocin (568 U) with 300 mL of 0.9% sodium chloride solution and adjusting the pH to 4.01 with 10× diluted acetic acid. The placebo spray (pH 4.01) was formulated with 0.9% sodium chloride solution and acetic acid, without the addition of oxytocin. The filtered and sterilized solutions were sealed in individual vials (1.5 mL each) and were frozen for storage. On the day of use, the vials were thawed and kept in a refrigerator at 4°C until required. A clinician prepared the nasal spray by transferring the oxytocin or the placebo from



**Fig 1. A consort diagram describing the participants in the study.**

doi:10.1371/journal.pone.0137514.g001

the vial into a nebulizer. The nebulizer was primed and given to the participants, who self-administered the nasal spray while being monitored by the clinician.

## Dynamic facial morphing task


The dynamic facial morphing task consisted of video clips showing computer-morphed faces. Neutral faces morphed into emotional faces showing sad, fearful, angry, or happy expressions with 50 gradual levels of each emotion, adhering to the experimental paradigm described by Joormann and Gotlib [35]. The task was formulated by one of the authors [36] and was conducted using MATLAB version R2010 (Mathworks, Natick, MA, USA) in conjunction with the Psychophysics Toolbox version 2.54 [37,38]. The facial stimuli were (12.7 × 16.9 cm) in size and were presented in colour in the middle of the screen with a grey background. The animated morphs were presented on a high-resolution 15-inch monitor, and the distance between the participant and the monitor was of about 60 cm.

The participants were asked to respond as quickly as possible with a keyboard stroke when they noticed any emotion in the face. The stimuli included faces of two men and two women taken from a database of Korean actors and actresses in their 20s to 40s [39]. Four practice trials and 16 experimental trials, which comprised four emotions from each of the four chosen actors/actresses, were conducted. The outcome was measured as the intensity level at which emotions were accurately identified (; emotional sensitivity).

In the task, a fixed '+' was initially shown for 500 ms in the centre of the screen, after which a neutral facial expression with 0% emotional intensity was shown for 500 ms. Progressive facial expressions of a 2% higher emotional intensity were shown at 500 ms intervals until the participant detected the presence of an emotion in the facial expression and reacted by pressing the space bar on the computer keyboard. The facial morphing was stopped, and the participant identified the emotion as 'sadness', 'fear', 'anger', or 'happiness' using the number keys '0', '1', '2', or '3', respectively, after which the next trial started immediately (Fig 2).

## Procedures

Each participant visited the laboratory twice for testing: once in the placebo condition and once in the oxytocin condition. Neither the participant nor the researchers involved were informed about the drug assignment of the day. The order in which to administer the placebo or the oxytocin to each participant was determined randomly (1:1) by the project coordinator,



**Fig 2. Dynamic facial morphing task.** In the task, a fixed '+' was initially shown for 500 ms in the centre of the screen, after which a neutral facial expression with 0% emotional intensity was shown for 500 ms. Progressive facial expressions of a 2% higher emotional intensity were shown at 500 ms intervals until the participant detected the presence of an emotion in the facial expression and reacted by pressing the space bar on the computer keyboard. The facial morphing was stopped, and the participant identified the emotion as 'sadness', 'fear', 'anger', or 'happiness' using the number keys '0', '1', '2', or '3', respectively, after which the next trial started immediately. The photograph is from a database published [39].

doi:10.1371/journal.pone.0137514.g002



who was not involved in conducting the experiment, using Microsoft Excel as follows: (1) The RAND function was used to generate a random number between 0 and 1, (2) If the generated number was smaller than 0.5, the participant was given the placebo first and (3) If the generated number was greater than or equal to 0.5, the participant was given oxytocin first.

The patients with eating disorders were provided with meal plans with fixed-size portions during the experimental period in order to not influence the drug's effect on calorie consumed during 24 hours. The patients with AN did not receive any direct support for eating during the 24 hours following the experiments on the inpatient ward. The patients with BN had meal plans focused on preventing bingeing and purging while supplying adequate energy to maintain a healthy weight, and they were instructed with a meal routine during the experimental period. Healthy controls were instructed to continue their routine diet during the 24 hours after the experiments.

The participants were tested in a private room at the Seoul Paik Hospital at 14:30. They were instructed to abstain from consuming alcohol or caffeine on the test day and to abstain from consuming food and drink (other than water) for 2 hours before the tests. Upon arrival, the participants completed baseline measures of physical symptoms, including abdominal, neurological, dermatological, and cardiac symptoms. The oxytocin and the placebo were self-administered intra-nasally under the supervision of a clinician, and the neuropsychological tasks were started 45 minutes later. A dose of 40 IU of oxytocin was chosen based on the results of a recent review that found the short-term use of intranasal oxytocin administered to male and female humans in dosages of up to 40 IU (per dose) to result in no detectable, subjective changes in a controlled research setting [40]. The nasal administration of oxytocin followed the guidelines established by Guastella et al. [41]. 90 min after the drug had been administered, the participants were asked to drink a carton of apple juice (Delmont, 190 mL per carton, 60 kcal). They were asked to consume as much of the juice as possible, as quickly as they were comfortable with, within a 20 min period. Finally, the participants completed an adverse symptom questionnaire. Over 24 hours following the experiment, the participants recorded a food diary to report every food and beverage items consumed, including the type, quantity, and approximate time and they sent back their diaries next day through their smartphones. A researcher who was blind to all the study procedures counted the calorie intake from the participant's diary. The second appointment was scheduled at least four days and at most seven days after the first appointment. The same procedure was repeated during the next visit, but the content of the spray differed according to the randomization plan.

## Statistical analysis

Sample size was calculated to detect two-way interaction effect using a repeated-measures ANOVA based on mixed effect model with 95% power and correlation among repeated measure = 0.5 across 3 groups. The results showed that 22 subjects would be needed with a medium effect ( $\Delta\eta^2 = 0.06$ ) and 129 subjects would be needed with a small effect ( $\Delta\eta^2 = 0.01$ ) per each group to detect significant interaction effect.

At the placebo condition, one-way ANOVAs were calculated for the amount of juice drank, calories consumed during 24 hours, and sensitivity to emotional recognition to examine the baseline differences among the diagnostic subtype (AN, BN, controls), as we valued the findings at the baseline.

We explored a mixed factor analysis (ANOVA) defining the following factors: 1) pre-post measure for drug (intra-factor); 2) drug (oxytocin-placebo) (intra-factor); and 3) diagnosis (AN-BN-controls) (inter-factor). As the pre-post measure was not significant on outcome measures of juice intake ( $p = 0.825$ ), calories consumed ( $p = 0.908$ ) and emotional intensity

( $p = 0.859$ ), we excluded carry-over effect of pre-post measure. Therefore, drug and diagnosis were included as factors in repeated measures ANOVA. For the food intake outcomes of the immediate juice intake and the calories consumed over 24 hours, we examined  $2(\text{drug}) \times 3(\text{diagnosis})$  repeated-measures ANOVAs and then made an independent analysis stratified by diagnosis to cope with low power due to small sample size.

As the level of difficulty of the morphing task was low, the main dependent variable we used was emotional intensity not accuracy. The sensitivity of emotional recognition was restricted to trials in which facial expressions were correctly identified. The emotional intensity was converted into the corrected emotional intensity which was calculated from the relative emotional intensity at the point of identification of an emotion.

The responses to the emotional stimuli were investigated via two way  $2(\text{drug}) \times 3(\text{diagnosis})$  repeated-measures analyses of variance (ANOVAs) on overall emotion, and then repeated-measures ANOVAs for each of the emotions (sad, fear, angry and happy) stratified by diagnosis were carried out. Post-hoc pairwise comparisons or simple main effect analyses were performed for significant main or interaction effects.

The data shown are presented as the mean and effect size (ES) if appropriate [Cohen's  $d$  for paired  $t$ -tests; partial eta squared ( $\Delta\eta^2$ ) for ANOVA]. Cohen's  $d$  was described as negligible ( $< 0.20$ ), small ( $< 0.50$ ), medium ( $\geq 0.50$  and  $< 0.8$ ), or large ( $\geq 0.8$ ) [42].  $\Delta\eta^2$  was described as small ( $< 0.01$ ), medium ( $< 0.059$ ), or large ( $< 0.138$ ) [43].  $P$ -values  $< 0.05$  were considered to be significant, and two-tailed tests were used. The statistical analyses were performed using SPSS 19.0 (IBM Inc., Chicago, IL, USA).

## Results

### Demographic and clinical characteristics

The demographic and clinical characteristics of the participants are shown in Table 1. The age ( $p = 0.754$ ) and IQ levels ( $p = 0.243$ ) did not differ among the diagnoses. The mean age of onset ( $p = 0.957$ ) and the mean duration of the illness ( $p = 0.243$ ) did not differ between patients with AN and patients with BN. Patients with AN and BN had higher levels of depression and anxiety than controls [all  $p < 0.001$ ,  $\text{BN} = \text{AN} > \text{HC}$  in post hoc Tukey]. The eating disorder psychopathology differed among the diagnoses [all  $p < 0.001$ : in post hoc Tukey,  $\text{BN} > \text{AN} > \text{HC}$  in global score and eating concern scale,  $\text{BN} > \text{AN} = \text{HC}$  in shape and weight concern scales,  $\text{AN} = \text{BN} > \text{HC}$  in restraint scale].

### Immediate consummatory behaviour of juice intake

Table 2 shows the results of immediate test meal of juice intake after receiving oxytocin or placebo. In the baseline condition, the amount of juice drank was significantly different among the groups [ $F(2,99) = 8.738$ ,  $p < 0.01$ ;  $\Delta\eta^2 = 0.150$ ]. The post hoc tests showed that both the patients with AN [ $p < 0.001$ ] and BN [ $p = 0.030$ ] drank a lower amount of juice than the controls did.

The two-way  $2(\text{drug}) \times 3(\text{diagnosis})$  repeated measures ANOVA showed a small effect of oxytocin [ $F(1,99) = 4.469$ ,  $p = 0.037$ ,  $\Delta\eta^2 = 0.043$ ] and large effect of diagnosis [ $F(2,99) = 11.600$ ,  $p < 0.001$ ,  $\Delta\eta^2 = 0.190$ ]. In the repeated measures ANOVAs stratified by diagnosis, there was no significant effect of oxytocin on juice intake in any diagnostic subtype.

### Caloric intake for 24-hours

Table 2 shows the caloric intake over a period of 24 hours after receiving oxytocin or placebo. In the baseline condition, the amount of calorie intake during 24 hours was different among



**Table 1. Demographic and clinical characteristics of healthy controls and the patients with anorexia nervosa and bulimia nervosa.**

Characteristics	AN (a, n = 35)	BN (b, n = 34)	HC (c, n = 33)	F(2,99)	p	Tukey HSD
Age, years	21.97(8.41)	23.03(5.17)	22.64(2.28)	0.283	0.754	NA
Age of onset, years	18.80(5.52)	18.74(4.38)	NA	$T_{(67)} = -0.054$	0.957	NA
Duration of illness, months	43.26(62.13)	57.59(34.72)	NA	$T_{(67)} = 1.178$	0.243	NA
WAIS, IQ	107.11(12.38)	106.62(11.91)	110.79(5.54)	1.581	0.211	NA
BMI	15.07(2.41)	20.24(2.45)	20.86(2.14)	63.334	<0.001	b = c>a
BDI	21.97(12.99)	18.29(10.94)	7.28(6.90)	16.788	<0.001	a = b>c
STAI						
State	55.85(13.62)	53.26(12.78)	43.44(11.27)	8.746	<0.001	a = b>c
Trait	54.94(13.29)	53.26(12.79)	43.75(11.16)	10.972	<0.001	a = b>c
EDE-Q						
Restraint	2.21(1.77)	2.32(1.78)	0.76(0.76)	10.927	<0.001	a = b>c
Eating Concern	1.88(1.69)	2.76(1.76)	0.56(0.77)	18.613	<0.001	b>a>c
Shape Concern	2.38(1.49)	3.56(1.82)	1.52(1.16)	15.343	<0.001	b>a = c
Weight Concern	2.73(1.45)	3.88(1.60)	2.21(1.29)	11.600	<0.001	b>a = c
Global	2.30(1.45)	3.13(1.48)	1.26(0.87)	17.413	<0.001	b>a>c

AN, anorexia nervosa; BN, bulimia nervosa; HC, healthy controls; WAIS: Wechsler Adult Intelligent Scale; IQ, Intelligent Quotient; BMI, body mass index; EDE-Q, Eating Disorder Examination Questionnaire; BDI, Beck Depression Inventory; STAI-State, Spielberger State and Trait Anxiety Inventory State score; STAI-Trait, Spielberger State and Trait Anxiety Inventory Trait score; NA, Not applicable

doi:10.1371/journal.pone.0137514.t001

the groups [ $F(2,99) = 6.581$ ,  $p = 0.002$ ,  $\Delta\eta^2 = 0.120$ ]. The post hoc tests showed that the patients with BN reported eating more calories than the patients with AN during the 24-hour period ( $p = 0.001$  for AN vs BN;  $p = 0.372$  for AN vs HC;  $p = 0.082$  for BN vs HC).

The two-way 2 (drug)  $\times$  3 (diagnosis) repeated-measures ANOVA showed a significant interaction between diagnosis and drug [ $F(2,99) = 4.736$ ,  $p = 0.011$ ,  $\Delta\eta^2 = 0.089$ ], but the effects of each of the factors were weak [for diagnosis,  $F(2,99) = 3.099$ ,  $p = 0.050$ ,  $\Delta\eta^2 = 0.060$ ; for drug,  $F(1,99) = 2.835$ ,  $p = 0.095$ ,  $\Delta\eta^2 = 0.028$ ]. In the repeated measures ANOVAs stratified by diagnosis, there was a significant effect of drug in patients with BN [ $F(1,33) = 6.389$ ,  $p = 0.016$ ,  $\Delta\eta^2 = 0.162$ ]. The patients with BN consumed fewer calories after receiving oxytocin ( $p = 0.016$ ,  $d = 0.560$ ). There was no significant effect of oxytocin either in healthy controls ( $p = 0.384$ ) or patients with AN ( $p = 0.145$ ).

## Sensitivity of emotional recognition

The thresholds for the facial emotion identification in the oxytocin condition are depicted in Table 3. In the baseline condition, there was no effect of diagnosis on the threshold of emotional recognition [ $F(2,99) = 0.662$ ,  $p = 0.518$ ,  $\Delta\eta^2 = 0.013$ ]. There was an effect of emotional type [ $F(3,97) = 171.584$ ,  $p < 0.001$ ,  $\Delta\eta^2 = 0.844$ ], in that happy emotion was easier to identify and angry emotion was less easy regardless of diagnostic subtype.

As the sample size was small, we made 2 (drug)  $\times$  3 (group) repeated-measures ANOVAs on overall emotion and then made an independent analysis stratified by diagnosis. The two-way 2 (drug)  $\times$  3 (group) repeated-measures ANOVAs demonstrated a main effect of drug [ $F(1,99) = 5.172$ ,  $p = 0.025$ ,  $\Delta\eta^2 = 0.053$ ] but no effect of group ( $p = 0.949$ ) (Fig 3).

In the stratified analysis by diagnosis, there was a small effect of drug in overall emotional recognition [ $F(1,32) = 4.530$ ,  $p = 0.041$ ,  $\Delta\eta^2 = 0.124$ ] in healthy controls. In the post hoc analysis, the sensitivity to overall emotional recognition was increased in the oxytocin condition in the healthy controls [ $t(32) = 2.128$ ,  $p = 0.041$ ,  $d = 0.311$ ]. In the following repeated-measures

**Table 2. The effect of oxytocin on immediate drinking juice and food intake for 24 hours in patients with anorexia nervosa and bulimia nervosa, and healthy university student.**

	AN (n = 35)				BN (n = 34)				HC (n = 33)				F	p	$\Delta r^2$
	Placebo	Oxytocin	t	p	Cohen's d	Placebo	Oxytocin	t	p	Cohen's d	Placebo	Oxytocin	t	p	Cohen's d
Juice drunk (ml)	92.03 (83.78)	97.49 (81.97)	-0.537	0.595	-0.082	118.35 (63.92)	131.50 (61.19)	-1.662	0.106	-0.197	160.91 (52.88)	172.82 (43.31)	-1.924	0.063	-0.179
Intake (kcal/day)	1988.55 (729.79)	2151.52 (873.33)	-1.492	0.145	-0.190	2757.84 (1047.65)	2277.60 (942.65)	2.528	0.016	0.560	2295.79 (808.18)	2179.48 (692.61)	0.882	0.384	0.136

AN, anorexia nervosa; BN, bulimia nervosa; HC, healthy controls.

Data are shown as mean (s.d.).

F: univariate ANOVA among 3 groups (AN, BN, HC) for placebo condition

t: paired t-test between placebo and oxytocin condition in each of the group

doi:10.1371/journal.pone.0137514.t002

**Table 3. Mean emotional intensity (%) of facial expressions made by patients with anorexia nervosa and bulimia nervosa, and by healthy controls.**

Emotion	AN (n = 35)				BN (n = 34)				HC (n = 33)				F	p	$\Delta\eta^2$
	placebo	oxytocin	t	p	Cohen's d	placebo	oxytocin	t	p	Cohen's d	placebo	oxytocin	t	p	Cohen's d
Total	40.56 (10.93)	40.85 (10.11)	0.238	0.814	-0.032	42.06 (8.83)	39.49 (7.40)	2.021	0.052	0.285	42.58 (8.22)	40.05 (8.05)	2.128	0.041	0.280
Sad	41.438 (11.285)	41.392 (10.895)	0.037	0.971	0.005	44.259 (12.729)	40.197 (7.781)	2.261	0.031	0.399	44.253 (8.504)	41.158 (8.939)	2.407	0.022	0.304
Fear	41.467 (10.455)	42.628 (10.714)	-0.885	0.383	-0.127	42.907 (7.143)	42.670 (8.380)	0.175	0.862	0.026	44.218 (8.813)	41.848 (8.968)	1.380	0.177	0.258
Angry	49.227 (13.727)	49.511 (12.644)	-0.147	0.884	-0.026	52.433 (9.130)	48.774 (10.182)	1.647	0.109	0.332	51.864 (9.756)	49.438 (9.774)	1.517	0.139	0.220
Happy	30.121 (12.474)	29.871 (12.896)	0.128	0.899	0.023	28.657 (12.288)	26.317 (8.109)	1.351	0.187	0.211	29.994 (10.697)	27.754 (9.098)	1.656	0.107	0.202
														0.146	0.864
														0.003	0.003


AN, anorexia nervosa; BN, bulimia nervosa; HC, healthy controls.

Data are shown as mean (s.d.).

F: univariate ANOVA among 3 groups (AN, BN, HC) for placebo treatment

t: paired t-test between placebo and oxytocin treatment in each group

doi:10.1371/journal.pone.0137514.t003



**Fig 3. Emotional intensity for accurate emotion identification of total emotion under oxytocin/placebo conditions.** Emotional intensity for accurate emotion identification of total emotion under oxytocin/placebo conditions. There was a main effect of drug [ $F(1,99) = 5.172$ ,  $p = 0.025$ ,  $\Delta\eta^2 = 0.053$ ], in which oxytocin increased the sensitivity of overall emotional recognition with a small effect size in the healthy women ( $d = 0.311$ ) and the patients with BN ( $d = 0.315$ ). AN, anorexia nervosa; BN, bulimia nervosa.

doi:10.1371/journal.pone.0137514.g003

ANOVAs for each of the emotions in the healthy controls, the effect of oxytocin was significant in sad emotion [ $F(1,32) = 5.792$ ,  $p = 0.022$ ,  $\Delta\eta^2 = 0.153$ ]. In the post hoc analysis, the sensitivity to sad emotion recognition was increased in the oxytocin condition in the healthy controls ( $p = 0.022$ ,  $d = 0.280$ ).

In the patients with BN, there was a tendency of drug effect on overall emotional recognition [ $F(1,33) = 3.461$ ,  $p = 0.072$ ,  $\Delta\eta^2 = 0.095$ ]. In the post hoc analysis, oxytocin had a tendency to increase the sensitivity to overall emotional recognition in the patients with BN [ $t(32) = 2.128$ ,  $p = 0.052$ ,  $d = 0.315$ ]. In the following repeated-measures ANOVAs for each of the emotions, the effect of oxytocin was significant in sad emotion in the patients with BN [ $F(1,33) = 5.144$ ,  $p = 0.031$ ,  $\Delta\eta^2 = 0.135$ ]. In the post hoc analysis, the sensitivity to sad emotion recognition was increased in the oxytocin condition in the patients with BN ( $p = 0.031$ ,  $d = 0.399$ ).

In the patients with AN, there was no effect of drug [ $F(1,34) = 0.348$ ,  $p = 0.559$ ,  $\Delta\eta^2 = 0.010$ ]. In the following repeated-measures ANOVAs for each of the emotions, there was no effect of oxytocin on the sensitivity to any of the emotion recognition in the patients with AN.

## Discussion

The aim of this study was to examine the impact of an intranasal oxytocin challenge test on two possible maintaining factors of eating disorders: problems in appetite control and social cognition. A single dose of intranasal oxytocin decreased caloric intake over 24 hours with a moderate effect in patients with BN. The immediate effect of oxytocin on intake was not significant in any of groups. Oxytocin increased emotion recognition particularly for sad expressions in patients with BN and in healthy women. No effects of oxytocin were found in either outcomes in patients with AN.

The most prominent finding in this study was that patients with BN consumed fewer calories over 24 hours after a single dose of intranasal oxytocin. The mechanisms underlying oxytocin's reduction of caloric intake, particularly in the BN group, have not yet been defined. Previous work found that oxytocin reduced hedonic eating in test meal conditions in men [25, 26]. Possible mechanism underpinning this effect may be by reducing emotional eating through reducing stress [44]. A study of anti-obesity effects of oxytocin in free living conditions in obese rats [12] suggested that oxytocin may also affect energy homeostasis [11], glucose metabolism and lipid metabolism [12]. The positive results of this proof of principle experimental study suggest that it may be useful to further examine the role of oxytocin in patients with BN.

In addition to the impact on appetite and metabolism, oxytocin has an impact on social cognition. We used dynamic facial expressions to investigate emotion recognition as these resemble naturalistic interpersonal situations better than static facial expressions. We found that oxytocin improved emotion recognition for dynamic facial expressions, particularly for negative emotions in patients with BN and in healthy women. This result is consistent with a previous study [45] in which administering oxytocin enhanced overall emotion recognition of faces in healthy humans. Our results are also consistent with a previous study that used a similar emotion recognition tasks with morphing faces, in which oxytocin increased the sensitivity to emotion recognition in healthy men [46]. It is interesting that intranasal oxytocin did not change emotion recognition sensitivity in patients with AN even though they had a relatively higher dose per body weight of oxytocin due to their low weight. We previously reported an increase in methylation in the MT2 region of the OXTR promoter in patients with AN [47]. Therefore, it is possible that this reduced functionality accounts for the limited overall effects of oxytocin observed in this group.

There are some limitations in this study that need to be considered. First, we used a self-reported diary to record food consumption during the 24 hours observation period which may have decreased the sensitivity of the outcome measure. Objective measures of caloric intake may be more precise. Second, the technology used to examine the eating behaviour was simple. A design including a "taste test", a covert method of measuring the consumption of high fat, sugar palatable foods, would have been of value to test the hypothesis that oxytocin has a specific effect on hedonic eating. Similarly it would have been of value to measure the trait of food addiction. Third, three patients with AN and 3 patients with BN were taking stable dose of fluoxetine in our study, which may have confounded the results. It is common in clinical practice that patients with eating disorders take antidepressants which makes it ecologically relevant to include them in this sample. Furthermore, an additional analysis found no differences in the outcome measures in the response of this subgroup to the non-medicated patients. Fourth, it is possible that a more challenging social cognition task, such as "reading the mind" task, may increase the sensitivity to find differences. Lastly, other minor limitation was that the sample size was too small to detect a small interaction effect with a repeated-measures ANOVA. Therefore, it is possible that we could not detect any interaction effect on emotional intensity with this sample size.

In conclusion, the impact of oxytocin on appetite and social cognition varied between people with AN and BN. A single dose of intranasal oxytocin decreased caloric intake over 24 hours in people with BN. Moreover, people with BN showed enhanced emotional sensitivity under oxytocin condition similar to healthy controls. Those effects of oxytocin were not found in patients with AN.

## Supporting Information

**S1 Dataset. Raw data for participants.**  
(SAV)

**S1 Text. Informed consent\_English.**  
(DOC)

**S2 Text. Informed consent\_Korean.**  
(DOC)

**S3 Text. Protocol\_English.**  
(DOC)

**S4 Text. Protocol\_Korean.**  
(DOC)

**S5 Text. Consort checklist.**  
(DOC)

## Author Contributions

Conceived and designed the experiments: YK JT. Performed the experiments: YK JK. Analyzed the data: YK JE. Contributed reagents/materials/analysis tools: JY. Wrote the paper: YK JT.

## References

1. Blevins JE, Ho JM (2013) Role of oxytocin signaling in the regulation of body weight. *Rev Endocr Metab Disord* 14: 311–329. doi: [10.1007/s11154-013-9260-x](https://doi.org/10.1007/s11154-013-9260-x) PMID: [24065622](https://pubmed.ncbi.nlm.nih.gov/24065622/)
2. Insel TR (2010) The Challenge of Translation in Social Neuroscience: A Review of Oxytocin, Vasopressin, and Affiliative Behavior. *Neuron* 65: 768–779. doi: [10.1016/j.neuron.2010.03.005](https://doi.org/10.1016/j.neuron.2010.03.005) PMID: [20346754](https://pubmed.ncbi.nlm.nih.gov/20346754/)
3. Maguire S, O'Dell A, Touyz L, Russell J (2013) Oxytocin and Anorexia Nervosa: a review of the emerging literature. *Eur Eat Disorders Rev* 21: 475–478.
4. Sabatier N, Leng G, Menzies J (2013) Oxytocin, feeding, and satiety. *Front Endocrinol* 35: 1–10.
5. Choleris E, Devidze N, Kavaliers M, Pfaff DW (2008) Steroidal/neuropeptide interactions in hypothalamus and amygdala related to social anxiety. In: Neumann ID, Landgraf R, editors. *Advances in Vasopressin and Oxytocin: From Genes to Behaviour to Disease*. pp. 291–303.
6. Demitrack MA, Lesem MD, Listwak SJ, Brandt HA, Jimerson DC, Gold PW (1990) CSF oxytocin in anorexia nervosa and bulimia nervosa: clinical and pathophysiologic considerations. *Am J Psychiatry* 147: 882–886. PMID: [2356873](https://pubmed.ncbi.nlm.nih.gov/2356873/)
7. Lawson EA, Donoho DA, Blum JL, Meenaghan EM, Misra M, Herzog DB, et al. (2011) Decreased nocturnal oxytocin levels in Anorexia Nervosa are associated with low bone mineral density and fat mass. *J Clin Psychiatry* 72: 1546–1551. doi: [10.4088/JCP.10m06617](https://doi.org/10.4088/JCP.10m06617) PMID: [21903023](https://pubmed.ncbi.nlm.nih.gov/21903023/)
8. Lawson EA, Holsen LM, Santin M, Meenaghan E, Eddy KT, Becker AE, et al. (2012) Oxytocin secretion is associated with severity of disordered eating psychopathology and insular cortex hypoactivation in Anorexia Nervosa. *J Clin Endocrinol Metabol* 97: E1898–E1908.
9. Kim YR, Kim CH, Cardi V, Eom JS, Seong Y, Treasure J (2014) Intranasal oxytocin attenuates attentional bias for eating and fat shape stimuli in patients with anorexia nervosa. *Psychoneuroendocrinology* 44: 133–142. doi: [10.1016/j.psyneuen.2014.02.019](https://doi.org/10.1016/j.psyneuen.2014.02.019) PMID: [24703429](https://pubmed.ncbi.nlm.nih.gov/24703429/)
10. Kim YR, Kim CH, Park JH, Pyo J, Treasure J (2014) The Impact of Intranasal Oxytocin on Attention to Social Emotional Stimuli in Patients with Anorexia Nervosa: A Double Blind within-Subject Cross-over Experiment. *Plos One* 9: e90721. doi: [10.1371/journal.pone.0090721](https://doi.org/10.1371/journal.pone.0090721) PMID: [24603863](https://pubmed.ncbi.nlm.nih.gov/24603863/)
11. Chaves VE, Tilelli CQ, Britob NA, Brito MN (2013) Role of oxytocin in energy metabolism. *Peptides* 45: 9–14. doi: [10.1016/j.peptides.2013.04.010](https://doi.org/10.1016/j.peptides.2013.04.010) PMID: [23628372](https://pubmed.ncbi.nlm.nih.gov/23628372/)
12. Deblon N, Veyrat-Durebex C, Bourgoin L, Caillon A, Bussier A-L, Petrosino S, et al. (2011) Mechanisms of the Anti-Obesity Effects of Oxytocin in Diet-Induced Obese Rats. *Plos One* 6.



13. Wu Z, Xu Y, Zhu Y, Sutton AK, Zhao R, Lowel B, et al. (2012) An obligate role of oxytocin neurons in diet induced energy expenditure. *PLoS One* 7: e45167. doi: [10.1371/journal.pone.0045167](https://doi.org/10.1371/journal.pone.0045167) PMID: [23028821](https://pubmed.ncbi.nlm.nih.gov/23028821/)
14. Leng G, Onaka T, Caquineau C, Sabatier N, Tobin VA, Takayanagi Y (2008) Oxytocin and appetite. *Prog Brain Res* 170: 137–151. doi: [10.1016/S0079-6123\(08\)00413-5](https://doi.org/10.1016/S0079-6123(08)00413-5) PMID: [18655879](https://pubmed.ncbi.nlm.nih.gov/18655879/)
15. Parker JA, Bloom SR (2012) Hypothalamic neuropeptides and the regulation of appetite. *Neuropharmacology* 63: 18–30. doi: [10.1016/j.neuropharm.2012.02.004](https://doi.org/10.1016/j.neuropharm.2012.02.004) PMID: [22369786](https://pubmed.ncbi.nlm.nih.gov/22369786/)
16. Olszewski PK, Klockars A, Schioth HB, Levine AS (2010) Oxytocin as feeding inhibitor Maintaining homeostasis in consummatory behavior. *Pharmacology Biochemistry and Behavior* 97: 47–54.
17. Amico JA, Vollmer RR, Cai HM, Miedlar JA, Rinaman L (2005) Enhanced initial and sustained intake of sucrose solution in mice with an oxytocin gene deletion. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology* 289: R1798–R1806.
18. Nishimori K, Takayanagi Y, Yoshida M, Kasahara Y, Young LJ, et al. (2008) New aspects of oxytocin receptor function revealed by knockout mice: sociosexual behaviour and control of energy balance. *Advances in Vasopressin and Oxytocin: From Genes to Behaviour to Disease* 170: 79–90.
19. Nishimori K, Young LJ, Guo QX, Wang ZX, Insel TR, et al. (1996) Oxytocin is required for nursing but is not essential for parturition or reproductive behavior. *Proceedings of the National Academy of Sciences of the United States of America* 93: 11699–11704. PMID: [8876199](https://pubmed.ncbi.nlm.nih.gov/8876199/)
20. Takayanagi Y, Kasahara Y, Onaka T, Takahashi N, Kawada T, et al. (2008) Oxytocin receptor-deficient mice developed late-onset obesity. *Neuroreport* 19: 951–955. doi: [10.1097/WNR.0b013e3283021ca9](https://doi.org/10.1097/WNR.0b013e3283021ca9) PMID: [18520999](https://pubmed.ncbi.nlm.nih.gov/18520999/)
21. Miedlar JA, Rinaman L, Vollmer RR, Amico JA (2007) Oxytocin gene deletion mice overconsume palatable sucrose solution but not palatable lipid emulsions. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology* 293: R1063–R1068.
22. Sclafani A, Rinaman L, Vollmer RR, Amico JA (2007) Oxytocin knockout mice demonstrate enhanced intake of sweet and nonsweet carbohydrate solutions. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology* 292: R1828–R1833.
23. Mitra A, Gosnell BA, Schioth HB, Grace MK, Klockars A, et al. (2010) Chronic sugar intake dampens feeding-related activity of neurons synthesizing a satiety mediator, oxytocin. *Peptides* 31: 1346–1352. doi: [10.1016/j.peptides.2010.04.005](https://doi.org/10.1016/j.peptides.2010.04.005) PMID: [20399242](https://pubmed.ncbi.nlm.nih.gov/20399242/)
24. Zhang G, Bai H, Zhang H, Dean C, Wu Q, Li J, et al. (2011) Neuropeptide exocytosis involving synaptotagmin-4 and oxytocin in hypothalamic programming of body weight and energy balance. *Neuron* 69: 523–535. doi: [10.1016/j.neuron.2010.12.036](https://doi.org/10.1016/j.neuron.2010.12.036) PMID: [21315262](https://pubmed.ncbi.nlm.nih.gov/21315262/)
25. Ott V, Finlayson G, Lehnert H, Heitmann B, Heinrichs M, Born J, et al. (2013) Oxytocin Reduces Reward-Driven Food Intake in Humans. *Diabetes* 62: 3418–3425. doi: [10.2337/db13-0663](https://doi.org/10.2337/db13-0663) PMID: [23835346](https://pubmed.ncbi.nlm.nih.gov/23835346/)
26. Lawson E, Marengi D, DeSanti R, Holmes T, Schoenfeld D, et al. (2015 in press) Oxytocin Reduces Caloric Intake in Men. *Obesity*.
27. Arcelus J, Haslam M, Farrow C, Meyer C (2013) The role of interpersonal functioning in the maintenance of eating psychopathology: A systematic review and testable model. *Clin Psychol Rev* 33: 156–167. doi: [10.1016/j.cpr.2012.10.009](https://doi.org/10.1016/j.cpr.2012.10.009) PMID: [23195616](https://pubmed.ncbi.nlm.nih.gov/23195616/)
28. Caglar-Nazali HP, Corfield F, Cardi V, Ambwani S, Leppanen J, Olabintan O, et al. (2014) A systematic review and meta-analysis of 'Systems for Social Processes' in eating disorders. *Neurosci Biobehav Rev* 42: 55–92. doi: [10.1016/j.neubiorev.2013.12.002](https://doi.org/10.1016/j.neubiorev.2013.12.002) PMID: [24333650](https://pubmed.ncbi.nlm.nih.gov/24333650/)
29. First MB, Spitzer RL, Gibbon M, Williams JBW (2002) *Structured Clinical Interview for DSM-IV-TR Axis I Disorders-Patient Edition (SCID-I/P)*. New York: Biometrics Research Department, New York State Psychiatric Institute.
30. Morgan JF, Reid F, Lacey JH (1999) The SCOFF questionnaire: assessment of a new screening tool for eating disorders. *British Medical Journal* 319: 1467–1468. PMID: [10582927](https://pubmed.ncbi.nlm.nih.gov/10582927/)
31. Fairburn C, Beglin S (1994) Assessment of eating disorders—Interview or self-report questionnaire. *Int J Eat Disord* 16: 363–370. PMID: [7866415](https://pubmed.ncbi.nlm.nih.gov/7866415/)
32. Beck AT, Ward CH, Mendelson M, Mock JE, Erbaugh J (1961) An inventory for measuring depression. *Arch Gen Psychiatry* 4: 561–571. PMID: [13688369](https://pubmed.ncbi.nlm.nih.gov/13688369/)
33. Spielberger C, Gorsuch R, Lushene R, Vagg P, Facobs G (1983) *Manual for the State-Trait Anxiety Inventory, STAI (Form Y)*. Palo Alto: Consulting Psychologists Press.
34. Ward LC (1990) Prediction of verbal, performance, and full scale IQs from the seven subsets of the WAIS-R. *J Clin Psychol* 46: 436–440. PMID: [2212047](https://pubmed.ncbi.nlm.nih.gov/2212047/)

35. Joormann J, Gotlib IH (2006) Is this happiness I see? Biases in the identification of emotional facial expressions in depression and social phobia. *J Abnorm Psychol* 115: 705–714. PMID: [17100528](#)
36. Yang J (2009) Perceptual and cognitive bias for facial emotion in social anxiety: Yonsei University Graduate School.
37. Brainard DH (1997) The psychophysics toolbox. *Spatial Vision* 10: 433–436. PMID: [9176952](#)
38. Pelli DG (1997) The videotoolbox software for visual psychophysics: Transforming numbers into movies. *Spat Vis* 10: 437–442. PMID: [9176953](#)
39. Yonsei University Cognitive Science Institute (1998) Development of facial expression and body gestures database. Seoul: Ministry of Education and Science.
40. MacDonald E, Dadds M, Brennan J, Williams K, Levy F, Cauch AJ (2011) A review of safety, side-effects and subjective reactions to intranasal oxytocin in human research. *Psychoneuroendocrinology* 36: 1114–1126. doi: [10.1016/j.psyneuen.2011.02.015](#) PMID: [21429671](#)
41. Guastella AJ, Hickie IB, McGuinness MM, Otis M, Woods EA, Disinger HM, et al. (2013) Recommendations for the standardisation of oxytocin nasal administration and guidelines for its reporting in human research. *Psychoneuroendocrinology* 38: 612–625. doi: [10.1016/j.psyneuen.2012.11.019](#) PMID: [23265311](#)
42. Cohen J (1992) A power primer. *Psychological Bulletin* 112: 155–159. PMID: [19565683](#)
43. Cohen J (1988) Statistical power analysis for the behavioral sciences. New Jersey: Lawrence Erlbaum Associated Inc.
44. Heinrichs M, Baumgartner T, Kirschbaum C, Ehler U (2003) Social support and oxytocin interact to suppress cortisol and subjective responses to psychosocial stress. *Biological Psychiatry* 54: 1389–1398. PMID: [14675803](#)
45. Shahrestani S, Kemp AH, Guastella AJ (2013) The Impact of a Single Administration of Intranasal Oxytocin on the Recognition of Basic Emotions in Humans: A Meta-Analysis. *Neuropsychopharmacology* 38: 1929–1936. doi: [10.1038/npp.2013.86](#) PMID: [23575742](#)
46. Lischke A, Berger C, Prehn K, Heinrichs M, Herpertz S, Domes G (2012) Intranasal oxytocin enhances emotion recognition from dynamic facial expressions and leaves eye-gaze unaffected. *Psychoneuroendocrinology* 37: 475–481. doi: [10.1016/j.psyneuen.2011.07.015](#) PMID: [21862223](#)
47. Kim YR, Kim JH, Kim MJ, Treasure J (2014) Differential Methylation of the Oxytocin Receptor Gene in Patients with Anorexia Nervosa: A Pilot Study. *PLoS ONE* 9: e90721. doi: [10.1371/journal.pone.0090721](#) PMID: [24603863](#)